

# **Resonance and Capture of Jupiter Comets**

**Shane Ross**

Martin Lo (JPL), Wang Sang Koon and Jerrold Marsden (Caltech)

**Control and Dynamical Systems  
California Institute of Technology**

US-European Celestial Mechanics Workshop, 3-7 July 2000


**[shane@cds.caltech.edu](mailto:shane@cds.caltech.edu)**

## ■ Acknowledgements

- H. Poincaré, J. Moser
- C. Conley, R. McGehee
- C. Simó, J. Llibre, R. Martínez
- E. Belbruno, B. Marsden, J. Miller
- G. Gómez, J. Masdemont
- K. Howell, R. Wilson and the Purdue group



## ■ Jupiter Comets: e.g., *Oterma*

- *Rapid transition* from **outside** to **inside** Jupiter's orbit.
- *Captured temporarily* by Jupiter during transition.
- **Exterior** (2:3 resonance). **Interior** (3:2 resonance).





## ■ Previous Work

- Belbruno/B. Marsden [1997]
- Lo/Ross [1997] : Comets pass by  $L_1$  &  $L_2$ .
  - In Sun-Jupiter *rotating frame*, comets follow ***stable & unstable invariant manifolds***.
- Works by Moser/Conley/McGehee. Llibre/Martinez/Simó [1985].




## ■ Flow Near $L_1$ and $L_2$

- *Energy value*  $> L_2$ : *Hill's region* has ***neck*** about  $L_1$  &  $L_2$ .
- Comet makes transition through *equilibrium region* necks.
- Four orbit types: periodic, **asymptotic**, **transit** & **nontransit**.




## Jupiter Comets Use Heteroclinic Connection

- **Heteroclinic connection** between  $L_1$  &  $L_2$  periodic orbits.
  - Link with homoclinic orbits to make *chain*.
- Comets follow dynamical *channels* for rapid transition.




## Jupiter Comets: Following Dynamical Channels

- Consider comet *Oterma* from 1910 to 1980.
  - Determine energy during transition.
  - Compute **homoclinic-heteroclinic chain**.
  - Overlay **chain** with *Oterma's* orbit (at right).




## Jupiter Comets: Rapid Transition Mechanism

- Rapid transition between exterior/interior via
  - stable & unstable manifold *tubes* which contain transit orbits.




- Jupiter region Poincaré section:
  - $L_2$  **unstable tube** intersects  $L_1$  **stable tube**.
  - Contains exterior  $\longrightarrow$  interior orbits.




## ■ Tubes and Resonance Transition


- Tubes are *transport mechanism* connecting interior and exterior Hill's regions.
- Connect **mean motion resonances**.
  - e.g., *Oterma's 2:3*— $\rightarrow$ *3:2* transition.
- Can construct *Oterma*-like transition.




- Schematic of tube location. Poincaré sections are **solid lines**.
  - **Stable** manifolds in **green**,
  - **Unstable** manifolds in **red**.




- Interior region Poincaré section.




- Look at Poincaré section of tubes in *exterior region*.
  - $L_2$  orbit **stable** & **unstable** manifolds.






- Exterior region Poincaré section.



- Transition resonances for *Oterma's* energy:
  - Interior: 3:2
  - Exterior: 2:3
- 2:3 → 3:2 connected via Jupiter region.




- Pick initial condition in **2:3**— $\rightarrow$ **3:2** channel.
  - Forward/backward integrate **Oterma**-like resonance transition.




## ■ Temporary Capture Around Jupiter

- Must come into Jupiter region via tubes for capture.
- Several revolutions possible before departing.




## ■ Collision with Jupiter

- Some portion of tube intersects Jupiter.
  - e.g., *Shoemaker-Levy 9* simulation




## ■ Further Work: Transport Throughout Solar System




## ■ More Information and References

- Koon, W.S., M.W. Lo, J.E. Marsden and S.D. Ross  
*Heteroclinic connections between periodic orbits and resonance transitions in celestial mechanics*,  
Chaos, vol. 10(2) [2000], pp. 427–469;
  - <http://www.cds.caltech.edu/~marsden/>
  - Click on “current issue” of  
<http://ojps.aip.org/chaos/>
- Email: [shane@cds.caltech.edu](mailto:shane@cds.caltech.edu)




- Look at Poincaré section of tubes in *Jupiter region*.

- $L_2$  orbit **unstable**  $\longrightarrow L_1$  orbit **stable** manifolds.





- Jupiter region Poincaré section:
  - $L_2$  **unstable tube** intersects  $L_1$  **stable tube**.



- Intersection region contains transit orbits:
  - exterior  $\longrightarrow$  interior


## Poincare Section in the Jupiter Region




## ■ Connection Between Interior/Exterior Resonances

- $\Delta$  contains orbits in  $2:3 \longrightarrow 3:2$  *transition*.
- Comets, e.g. *Oterma*, pass through such regions.

**Interior Region**



**Jupiter Region**



**Exterior Region**

